

Retraction

TAF_{II}s Mediate Activation of Transcription in the *Drosophila* Embryo

We previously reported that transcriptional activation of *hunchback* (*hb*) and *huckebein* (*hkb*) by Bicoid in the *Drosophila* embryo is impaired by mutations in TAF_{II}110 and TAF_{II}60. This conclusion was based on in situ staining of wild-type versus mutant embryos (Figures 5 and 6 of Sauer et al., Cell 87[7], 1271–1284, 1996). However, we have come to realize that, due to technical problems with our previous embryo staining procedures, the in vivo results in Figures 5 and 6 are incorrect and those in Figures 7–9 are uncertain. Recent data by J. Zhou and R. T. indicate that mutations in TAF_{II}60 (Figure 1 below) and TAF_{II}110 (data not shown) have no detectable effect on *hb* expression. However, the isolation and molecular characterization of these TAF_{II} mutants (Figure 1–3, Sauer et al., 1996) is correct and recent studies (see Zhou et al., 1998) suggest that these TAF_{II} mutations affect Dorsal-mediated activation of *snail* and *twist* in the *Drosophila* embryo. We thank Gary Struhl, Claude Desplan, and Mike Levine for alerting us to problems in our previous embryo staining experiments and Jumin Zhou for the results in Figure 1.

Frank Sauer,* David A. Wassarman,† Gerald M. Rubin,‡ and Robert Tjian‡

*ZMBH

Im Neunheimer Feld 282
D-69120 Heidelberg, Germany

†National Institute of Child Health
and Human Development
Cell Biology and Metabolism Branch
Bethesda, Maryland 20892

‡Department of Molecular and Cell Biology
Howard Hughes Medical Institute
University of California at Berkeley
Berkeley, California 94720

References

- Lehmann, R., and Nüsslein-Volhard, C. (1987). *hunchback*, a gene required for segmentation of an anterior and posterior region of the *Drosophila* embryo. Dev. Biol. 119, 402–417.
- Zhou, J., Zwicker, J., Szymanski, P., Levine, M., and Tjian, R. (1998). TAF_{II} mutations disrupt Dorsal activation in the *Drosophila* embryo. Proc. Natl. Acad. Sci. USA, in press.




Figure 1. *hunchback* (*hb*) Expression in TAF_{II}60^Y Mutant Embryos

Stage 5 embryos were collected from the mating of *hb*¹⁴/*P*[*Kr-lacZ*] heterozygous males and either wild-type (A and B) or TAF_{II}60^Y/*+* heterozygous (C and D) females. The embryos (2.5 hr postfertilization) were hybridized with a mixture of digoxigenin-labeled *hb* and *lacZ* antisense RNA probes; they are oriented with anterior to the left and dorsal up.

(A) *hb* expression in a *+/P*[*Kr-lacZ*] embryo that contains two normal cop-

ies of the *hb*⁺ gene, as indicated by the expression of the *Kr-lacZ* reporter gene in central regions (long arrow). Normal patterns of *hb* expression can be detected in both anterior and posterior regions of the embryo.

(B) *hb* expression in an embryo that contains only one normal copy of the *hb*⁺ gene, as indicated by the absence of *lacZ* expression (small arrow) in the *Kr* domain. *hb* transcripts are detected in both anterior and posterior regions. However, there is an ~2-fold reduction in the levels of *hb* transcripts (compare with A) since the *hb*¹⁴ null mutation used in this analysis does not encode a stable transcript (Lehmann and Nüsslein-Volhard, 1987). (C) Embryo of genotype *+/P*[*Kr-lacZ*] that was derived from a TAF_{II}60^Y/*+* heterozygous female. The reduction in the maternal dose of the wild-type TAF_{II}60 product does not alter the limits or levels of the *hb* expression pattern (compare with A).

(D) Embryo of genotype *hb*¹⁴/*+* that was derived from a TAF_{II}60^Y/*+* heterozygous female. The simultaneous reduction in *hb*⁺ and TAF_{II}60⁺ gene does not alter the *hb* expression pattern beyond the 2-fold reduction in staining due to the *hb*¹⁴ null mutation (compare with A and B).